

PE Rabbit anti-Human CD8 PolymAb®

Catalog No.: A28209PM

Basic Information

Observed MW

Refer to figures

Calculated MW

26 kDa/24 kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

Conjugate

PE. Ex:565nm. Em:574nm.

Background

The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a coreceptor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell in the context of class I MHC molecules. The coreceptor functions as either a homodimer composed of two alpha chains or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 alpha chain. Multiple transcript variants encoding different isoforms have been found for this gene. The major protein isoforms of this gene differ by the presence or absence of a transmembrane domain and thus differ in being a membrane-anchored or secreted protein.

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

925/926

Swiss Prot

P01732/P10966

Immunogen

This information is considered to be commercially sensitive.

Synonyms

CD8; p32; Leu2; IMD116; CD8alpha; LY3; P37; LEU2; LYT3; Ly-3; CD8B1; CD8beta.

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1X10⁶ Human PBMC were surface-stained with ABflo® 488 Rabbit anti-Human/Monkey CD3 mAb (A26283,5 µl/Test) and PE Rabbit IgG isotype control (A24172,5 µl/Test, left) or PE Rabbit anti-Human CD8 PolymAb® (A28209PM,5 µl/Test, right). Cells in the lymphocyte gate were used for analysis.