

COX2/PTGS2 Rabbit pAb

Catalog No.: A28291

Basic Information

Observed MW

74 kDa

Calculated MW

69 kDa

Category

Primary antibody

Applications

WB, IF/ICC, ELISA

Cross-Reactivity

Mouse

Background

This gene encodes an enzyme that is a member of the prostaglandin G/H synthase family. The encoded protein converts arachidonic acid to prostaglandin endoperoxide H₂ which is a key enzymatic step in prostaglandin biosynthesis. This gene is the inducible prostaglandin G/H synthase family member that is upregulated during inflammation. Aberrant regulation of this gene is associated with cancer progression in several tissues and an increased risk of cardiovascular events. This gene is the target of many non-steroidal anti-inflammatory drugs.

Recommended Dilutions

WB 1:1000 - 1:3000

IF/ICC 1:100 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

19225

Swiss Prot

Q05769

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

COX2; Cox-2; PES-2; PHS-2; Pgbs2; TIS10; PGHS-2; PHS II; gripghs

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from RAW 264.7 cells using COX2/PTGS2 Rabbit pAb (A28291) at 1:3000 dilution incubated overnight at 4°C. RAW 264.7 cells were treated with LPS (1 µg/mL) for 24 hours.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60 s.



Immunofluorescence analysis of RAW 264.7 cells (treated with LPS) and RAW 264.7 cells (untreated) using COX2/PTGS2 Rabbit pAb (A28291) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.