

XBP-1s Rabbit mAb

Catalog No.: A28350 **Recombinant**

Basic Information

Observed MW

56 kDa

Calculated MW

40 kDa

Category

Primary antibody

Applications

WB,ChIP,ELISA

Cross-Reactivity

Human, Mouse

CloneNo number

ARC78803

Background

This gene encodes a transcription factor that regulates MHC class II genes by binding to a promoter element referred to as an X box. This gene product is a bZIP protein, which was also identified as a cellular transcription factor that binds to an enhancer in the promoter of the T cell leukemia virus type 1 promoter. It may increase expression of viral proteins by acting as the DNA binding partner of a viral transactivator. It has been found that upon accumulation of unfolded proteins in the endoplasmic reticulum (ER), the mRNA of this gene is processed to an active form by an unconventional splicing mechanism that is mediated by the endonuclease inositol-requiring enzyme 1 (IRE1). The resulting loss of 26 nt from the spliced mRNA causes a frame-shift and an isoform XBP1(S), which is the functionally active transcription factor. The isoform encoded by the unspliced mRNA, XBP1(U), is constitutively expressed, and thought to function as a negative feedback regulator of XBP1(S), which shuts off transcription of target genes during the recovery phase of ER stress. A pseudogene of XBP1 has been identified and localized to chromosome 5.

Recommended Dilutions

WB 1:2000 - 1:5000

ChIP 1-3µg antibody for
10µg-15µg of Chromatin

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions ($\geq 1:10000$) a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID

7494

Swiss Prot

P17861-2

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

XBP2; TREB5; XBP-1; TREB-5

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Contact


 | 400-999-6126

 | cn.market@abclonal.com.cn
 | www.abclonal.com.cn


Validation Data



Western blot analysis of lysates from HeLa cells using XBP-1s Rabbit mAb (A28350) at 1:5000 dilution incubated overnight at 4°C. HeLa cells were treated with Tunicamycin (20 µg/mL) for 4 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 30 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 45 s.



Chromatin immunoprecipitation was performed with 15 µg of cross-linked chromatin from C2C12 cells treated with tunicamycin (2 µg/ml, 8h), using 1 µg of XBP-1s Rabbit mAb (A28350) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.



Chromatin immunoprecipitation was performed with 15 µg of cross-linked chromatin from HeLa cells treated with tunicamycin (2 µg/ml, 7h), using 3 µg of XBP-1s Rabbit mAb (A28350) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.