APC Rabbit anti-Human CD271/NGFR mAb

ABclonal www.abclonal.com

Catalog No.: A28359

Basic Information

Observed MW

Calculated MW

35kDa/45kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC57800

Conjugate

APC. Ex:650nm. Em:660nm.

Background

Nerve growth factor receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region, a single transmembrane domain, and a 155-amino acid cytoplasmic domain. The cysteine-rich region contains the nerve growth factor binding domain.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 4804 **Swiss Prot**

P08138

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data









Flow cytometry:1X10^6 K-562 cells (negative control,left) and SK-N-MC cells (right) were surface-stained with APC Rabbit anti-Human CD271/NGFR mAb(A28359,5 $\mu | \text{Test,orange line})$ or APC Rabbit IgG isotype control (A24173,5 $\mu | \text{Test,blue line})$. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 SK-N-MC cells were surface-stained with APC Rabbit IgG isotype control (A24173,5 μ I/Test,Ieft) or APC Rabbit anti-Human CD271/NGFR mAb(A28359,5 μ I/Test,right).