PE/Cyanine7 Rabbit anti-Mouse Ki67 mAb

www.abclonal.com

ABclonal

Catalog No.: A28387

Basic Information

Observed MW

Calculated MW

351 kDa

Category

Primary antibody

Applications

FC (intra)

Cross-Reactivity

Mouse

CloneNo number

ARC69121

Conjugate

PE-Cy7. Ex:565nm. Em:778nm.

Background

The nuclear protein Ki-67 was initially identified by the monoclonal antibody Ki-67, which was generated through immunization of mice with nuclei from the L428 Hodgkin's lymphoma cell line. The Ki-67 protein plays a crucial role in ribosomal RNA transcription and cell proliferation. Its expression is detected during the G1, S, G2, and M phases of the cell cycle, while it is absent in the G0 phase. Ki-67 is strongly expressed in proliferating cells and has been reported as a prognostic marker for various types of tumors.

Recommended Dilutions

FC (intra)

≤0.25 µg per million cells in 100 µl volume

Immunogen Information

Gene ID

Swiss Prot

17345

E9PVX6

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Ki67; Ki-67; D630048A14Rik

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry:1X10^6 C57BL/6 mouse splenocytes (treated with ConA for 3day) were intracellularly-stained with PE/Cyanine7 Rabbit IgG isotype control (0.25µg,left) or PE/Cyanine7 Rabbit anti-Mouse Ki67 mAb (A28387,0.25µg,right).