

# APC Rabbit anti-NHP CD45 mAb

Catalog No.: A28472

## Basic Information

### Observed MW

### Calculated MW

148 kDa

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

Cynomolgus, Rhesus

### CloneNo number

ARC79398

### Conjugate

APC. Ex:650nm. Em:660nm.

## Background

CD45 is a receptor-type protein tyrosine phosphatase, also known as Ly-5 or the leukocyte common antigen. It functions primarily by regulating the activation of Src family protein tyrosine kinases, such as Lck and Fyn, thereby initiating T cell receptor signaling. Deficiency in CD45 results in dysfunctional T and B lymphocytes, leading to severe combined immunodeficiency. Furthermore, CD45 plays a significant role in the pathogenesis of autoimmune diseases, cancer, and infectious diseases, including fungal infections.

## Recommended Dilutions

FC 5  $\mu$ l per  $10^6$  cells in  
100  $\mu$ l volume

## Immunogen Information

### Gene ID

102116047

### Swiss Prot

A0A2K5VQ77

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

Pan Leukocyte; NHP-specific; PTPRC; LCA; L-CA; Leukocyte Common Ag

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

## Validation Data



Flow cytometry:  $1 \times 10^6$  Human peripheral blood mononuclear cells (negative control, left) and Cynomolgus peripheral blood mononuclear cells (right) were surface-stained with APC Rabbit anti-NHP CD45 mAb (A28472, 5  $\mu$ l/Test).

Flow cytometry:  $1 \times 10^6$  Rhesus PBMC were surface-stained with APC Rabbit IgG isotype control (A24173, 5  $\mu$ l/Test, left) or APC Rabbit anti-NHP CD45 mAb (A28472, 5  $\mu$ l/Test, right). Cells in the lymphocyte gate were used for analysis.