# [KO Validated] p53 Rabbit mAb

Catalog No.: A28493 KO Validated Recombinant



### **Basic Information**

#### **Observed MW**

53 kDa

#### **Calculated MW**

24-44 kDa

### Category

Primary antibody

#### **Applications**

WB,IP,IF/ICC,IHC-P,ChIP,ELISA

#### **Cross-Reactivity**

Human

#### CloneNo number

ARC74241

### **Background**

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants.

### **Recommended Dilutions**

**WB** 1:2000 - 1:5000

**IF/ICC** 1:200 - 1:1000

IHC-P 1:400 -1600

**IP** 0.2 μg-4 μg antibody for

200 μg-400 μg extracts

of whole cells

**ChIP** 2μg antibody for

5μg-10μg of Chromatin

**ELISA** Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay
requirements.For highratio antibody dilutions
(≥1:10000)a sequential
dilution method is
strongly recommended

to ensure measurement

accuracy.

### Immunogen Information

**Gene ID**7157

Swiss Prot
P04637

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

P53; BCC7; LFS1; BMFS5; TRP53

### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification


#### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Contact

2	400-999-6126
$\bowtie$	cn.market@abclonal.com.cr
•	www.abclonal.com.cr



Immunoprecipitation of p53 from 300  $\mu$ g extracts of 293T cells was performed using 0.2  $\mu$ g of [KO Validated] p53 Rabbit mAb(A28493). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using [KO Validated] p53 Rabbit mAb(A28493) at a dilution of 1:10000.




Western blot analysis of lysates from wild type (WT) and p53 knockout (KO) 293T cells using [KO Validated] p53 Rabbit mAb (A28493) at 1:5000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45 s.




Western blot analysis of lysates from MCF7 cells using [KO Validated] p53 Rabbit mAb (A28493) at 1:5000 dilution incubated overnight at 4°C. MCF7 cells were treated with doxorubicin(0.5  $\mu$ M) at 37°C for 24 hours, MG-132(5  $\mu$ M) at 37°C for 6 hours by co-treatment.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 30 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 45 s.




Chromatin immunoprecipitation was performed with 15  $\mu$ g of cross-linked chromatin from HeLa cells and HeLa cells were treated with UV (100 mJ/cm²) at room temperature and recovered for 2 hours, using 2  $\mu$ g of [KO Validated] p53 Rabbit mAb (A28493) and Rabbit IgG isotype control(AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.



Confocal imaging of HT-29 cells using [KO Validated] p53 Rabbit mAb (A28493, dilution 1:600) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with  $\alpha$ -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunohistochemistry analysis of paraffinembedded Human ovarian serous carcinoma tissue using [KO Validated] p53 Rabbit mAb (A28493) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human colon carcinoma tissue using [KO Validated] p53 Rabbit mAb (A28493) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.