Leader in Biomolecular Solutions for Life Science

www.abclonal.com

ABflo® 488 anti-Mouse CD45.1 Antibody

Catalog No.: A28522

Basic Information

Observed MW

Calculated MW
130 kDa/140 kDa/144 kDa

Category
Primary antibody

Applications
FC

Cross-Reactivity
Mouse

CloneNo number
AMC0904

Conjugate
ABflo® 488. Ex:491nm. Em:516nm.

Background

Enables several functions, including heparan sulfate proteoglycan binding activity; heparin binding activity; and protein tyrosine phosphatase activity. Involved in several processes, including lymphocyte differentiation; positive regulation of macromolecule metabolic process; and regulation of signal transduction. Acts upstream of or within several processes, including lymphocyte differentiation; positive regulation of lymphocyte activation; and regulation of protein phosphorylation. Located in external side of plasma membrane; focal adhesion; and membrane raft. Is expressed in several structures, including 3rd branchial arch; alimentary system; cardiovascular system; hemolymphoid system; and placenta. Used to study systemic lupus erythematosus. Human ortholog(s) of this gene implicated in hepatitis C; multiple sclerosis; severe combined immunodeficiency; and severe combined immunodeficiency, autosomal recessive, T cell-negative, B cell-positive, Nk cell-positive. Orthologous to human PTPRC (protein tyrosine phosphatase receptor type C).

Recommended Dilutions

FC	1 μg per million cells in 100 μl volume

Immunogen Information

Gene ID
19264

Swiss Prot
P06800

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
loc; B220; Cd45; L-CA; Ly-5; T200; CD45R; Lyt-4

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information

Source
Mouse

Isotype
Mouse (A.SW) IgG2a, κ

Purification
Affinity purification

Storage
Store at -20℃