

Anti-Monkey CD117/c-Kit mAb

Catalog No.: A28853

Basic Information

Observed MW**Calculated MW**

110 kDa

Category

Primary antibody

Applications

IHC-P

Cross-Reactivity

Cynomolgus monkey

CloneNo number

ARC5250-01

Background

This gene encodes a receptor tyrosine kinase. This gene was initially identified as a homolog of the feline sarcoma viral oncogene v-kit and is often referred to as proto-oncogene c-Kit. The canonical form of this glycosylated transmembrane protein has an N-terminal extracellular region with five immunoglobulin-like domains, a transmembrane region, and an intracellular tyrosine kinase domain at the C-terminus. Upon activation by its cytokine ligand, stem cell factor (SCF), this protein phosphorylates multiple intracellular proteins that play a role in the proliferation, differentiation, migration and apoptosis of many cell types and thereby plays an important role in hematopoiesis, stem cell maintenance, gametogenesis, melanogenesis, and in mast cell development, migration and function. This protein can be a membrane-bound or soluble protein. Mutations in this gene are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous leukemia, and piebaldism. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

IHC-P 1:10000 - 1:40000

Immunogen Information

Gene ID

3815

Swiss Prot

P10721

Immunogen

This information is considered to be commercially sensitive.

Synonyms

PBT; SCFR; C-Kit; CD117; MASTC

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Cynomolgus monkey colon tissue using Anti-Monkey CD117/c-Kit mAb (A28853) at a dilution of 1:18000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Cynomolgus monkey kidney tissue using Anti-Monkey CD117/c-Kit mAb (A28853) at a dilution of 1:18000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.