

Neurofilament/NF-L Rabbit mAb

Catalog No.: A29439 **Recombinant**

Basic Information

Observed MW

70 kDa

Calculated MW

62 kDa

Category

Primary antibody

Applications

WB,IF/ICC,IF-F,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC4143

Background

Neurofilaments are type IV intermediate filament heteropolymers composed of light, medium, and heavy chains. Neurofilaments comprise the axoskeleton and they functionally maintain the neuronal caliber. They may also play a role in intracellular transport to axons and dendrites. This gene encodes the light chain neurofilament protein. Mutations in this gene cause Charcot-Marie-Tooth disease types 1F (CMT1F) and 2E (CMT2E), disorders of the peripheral nervous system that are characterized by distinct neuropathies. A pseudogene has been identified on chromosome Y.

Recommended Dilutions

WB 1:100000 - 1:400000**IF/ICC** 1:2000 - 1:8000**IF-F** 1:2000 - 1:8000**IHC-P** 1:2000 - 1:8000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (≥1:10000) a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID

4747

Swiss Prot

P07196

Immunogen

This information is considered to be commercially sensitive.

Synonyms

NFL; NF-L; NF68; CMT1F; CMT2E; CMTDIG; PPP1R110

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Western blot analysis of various lysates using Neurofilament/NF-L Rabbit mAb (A29439) at 1:200000 dilution incubated overnight at 4°C.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Negative control (NC): HeLa.
 Exposure time: 5 s.




Confocal imaging of PC-3 cells using Neurofilament/NF-L Rabbit mAb (A29439, dilution 1:4000) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of 293T cells using Neurofilament/NF-L Rabbit mAb (A29439, dilution 1:4000) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of frozen sections of Rat brain tissue using Neurofilament/NF-L Rabbit mAb (A29439, dilution 1:4000) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of frozen sections of Mouse brain tissue using Neurofilament/NF-L Rabbit mAb (A29439, dilution 1:4000) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffin-embedded Human brain tissue (left, Positive control) and Human liver tissue (right, Negative control), using Neurofilament/NF-L Rabbit mAb (A29439) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human brain tissue using Neurofilament/NF-L Rabbit mAb (A29439) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using Neurofilament/NF-L Rabbit mAb (A29439) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using Neurofilament/NF-L Rabbit mAb (A29439) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.