

ABflo® 647 Rabbit anti-Whitlow/218 Linker mAb

Catalog No.: A29518

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC

Cross-Reactivity

Species independent

CloneNo number

ARC84338

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

The Whitlow/218 peptide is a commonly used linker that connects the variable heavy (VH) domain and variable light (VL) domain of single-chain variable fragments (scFvs) and chimeric antigen receptors (CARs) that utilize an extracellular domain scFv for target antigen recognition. The Whitlow/218 linker affords the scFv with enhanced resistance to proteolysis and aggregation.

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 293T cells (negative control, left) and 293T cells engineered to express an scFv-based anti-CD19 CAR containing a Whitlow/218 linker (right) were surface-stained with ABflo® 647 Rabbit anti-Whitlow/218 Linker mAb (A29518, 0.25 μ g, orange line) or ABflo® 647 Rabbit IgG isotype control (A22070, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).