Leader in Biomolecular Solutions for Life Science

www.abclonal.com

NLRC4 Rabbit mAb

Catalog No.: A30149 Recombinant

Basic Information

Observed MW
110 kDa (C-terminal protein)

Calculated MW
116 kDa/11 kDa/18 kDa/41 kDa

Category
Primary antibody

Applications
WB,IF-P,IHC-P

Cross-Reactivity
Human

CloneNo number
ARC85313

Background

This gene encodes a member of the caspase recruitment domain-containing NLR family. Family members play essential roles in innate immune response to a wide range of pathogenic organisms, tissue damage and other cellular stresses. Mutations in this gene result in autoinflammation with infantile enterocolitis. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB	1:5000 - 1:10000
IF-P	1:200 -1:400
IHC-P	1:200 - 1:800

Immunogen Information

Gene ID
58484

Swiss Prot
Q9NPP4

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
CLAN; IPAF; AIFEC; CLAN1; CLANA; CLANB; CLANC; CLAND; FCAS4; CARD12; CLR2.1

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Validation Data




Western blot analysis of lysates from wild type (WT) and 293T cells transfected with NLRC4 using NLRC4 Rabbit mAb (A30149) at 1:5000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 20 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1 s.




Confocal imaging of paraffin-embedded Human liver tissue using NLRC4 Rabbit mAb (A30149, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using NLRC4 Rabbit mAb (A30149) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using NLRC4 Rabbit mAb (A30149) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human spleen tissue using NLRC4 Rabbit mAb (A30149) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using NLRC4 Rabbit mAb (A30149) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.