

# MYSM1 Rabbit pAb

Catalog No.: A3102

## Basic Information

### Observed MW

105kDa

### Calculated MW

95kDa

### Category

Primary antibody

### Applications

WB,ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

Enables histone binding activity; peptidase activity; and transcription coactivator activity. Involved in several processes, including chromatin remodeling; monoubiquitinated histone H2A deubiquitination; and positive regulation of transcription by RNA polymerase II. Located in nucleolus and nucleoplasm. Part of protein-containing complex. Implicated in diabetic retinopathy.

## Recommended Dilutions

**WB** 1:500 - 1:2000

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

114803

### Swiss Prot

Q5VVJ2

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

2ADUB; BMFS4; 2A-DUB; MYSM1

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of various lysates using MYSM1 Rabbit pAb (A3102) at 1:3000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.