DNMT3A Rabbit pAb

Catalog No.: A3169 5 Publications



Basic Information

Observed MW

95kDa

Calculated MW

102kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

CpG methylation is an epigenetic modification that is important for embryonic development, imprinting, and X-chromosome inactivation. Studies in mice have demonstrated that DNA methylation is required for mammalian development. This gene encodes a DNA methyltransferase that is thought to function in de novo methylation, rather than maintenance methylation. The protein localizes to the cytoplasm and nucleus and its expression is developmentally regulated.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:20

ELISA Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 1788 Q9Y6K1

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

TBRS; HESJAS; DNMT3A2; M.HsallIA; DNMT3A

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	Τ	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using DNMT3A Rabbit pAb (A3169).


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.




Western blot analysis of various lysates using DNMT3A Rabbit pAb (A3169) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates / proteins: 25 μ g per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of A549 cells using DNMT3A Rabbit pAb (A3169). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution.