

USP7/HAUSP Rabbit mAb

Catalog No.: A3448 **Recombinant** **2 Publications**

Basic Information

Observed MW

140kDa

Calculated MW

128kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC0777

Background

The protein encoded by this gene belongs to the peptidase C19 family, which includes ubiquitinyl hydrolases. This protein deubiquitinates target proteins such as p53 (a tumor suppressor protein) and WASH (essential for endosomal protein recycling), and regulates their activities by counteracting the opposing ubiquitin ligase activity of proteins such as HDM2 and TRIM27, involved in the respective process. Mutations in this gene have been implicated in a neurodevelopmental disorder.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200

Immunogen Information

Gene ID

7874

Swiss Prot

Q93009

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 100-200 of human USP7/HAUSP (Q93009).

Synonyms

TEF1; HAUSP; HAFOUS; DEL16P13.2; C16DELp13.2; USP7/HAUSP

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using USP7/HAUSP Rabbit mAb (A3448) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.




Immunohistochemistry analysis of USP7/HAUSP in paraffin-embedded human colon carcinoma tissue using USP7/HAUSP Rabbit mAb (A3448) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of USP7/HAUSP in paraffin-embedded human colon tissue using USP7/HAUSP Rabbit mAb (A3448) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of USP7/HAUSP in paraffin-embedded human thyroid cancer tissue using USP7/HAUSP Rabbit mAb (A3448) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of USP7/HAUSP in paraffin-embedded mouse brain tissue using USP7/HAUSP Rabbit mAb (A3448) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of USP7/HAUSP in paraffin-embedded mouse testis tissue using USP7/HAUSP Rabbit mAb (A3448) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of USP7/HAUSP in paraffin-embedded rat brain tissue using USP7/HAUSP Rabbit mAb (A3448) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of USP7/HAUSP in paraffin-embedded rat colon tissue using USP7/HAUSP Rabbit mAb (A3448) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of USP7/HAUSP in paraffin-embedded rat kidney tissue using USP7/HAUSP Rabbit mAb (A3448) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.