# **RhoGAP Rabbit pAb**

Catalog No.: A3587



### **Basic Information**

#### **Observed MW**

185kDa

### **Calculated MW**

172kDa

### Category

Primary antibody

### **Applications**

ELISA,WB,IF/ICC

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Rho GTPase activating protein 5 negatively regulates RHO GTPases, a family which may mediate cytoskeleton changes by stimulating the hydrolysis of bound GTP. Two transcript variants encoding different isoforms have been found for this gene.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

# **Immunogen Information**

**Gene ID**Swiss Prot
394
Q13017

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1292-1501 of human RhoGAP (NP\_001164.2).

### **Synonyms**

GFI2; p190-B; RhoGAP5; p190BRhoGAP; RhoGAP

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{a}$	ī	www.ahclonal.com.cn


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.




Western blot analysis of extracts of various cell lines, using RhoGAP antibody (A3587) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunofluorescence analysis of HeLa cells using RhoGAP antibody (A3587) at dilution of 1:100. Blue: DAPI for nuclear staining.