AMFR Rabbit pAb

Catalog No.: A3731 1 Publications



Basic Information

Observed MW

72kDa

Calculated MW

73kDa

Category

Primary antibody

Applications

WB,ELISA

Cross-Reactivity

Human

Background

This locus encodes a glycosylated transmembrane receptor. Its ligand, autocrine motility factor, is a tumor motility-stimulating protein secreted by tumor cells. The encoded receptor is also a member of the E3 ubiquitin ligase family of proteins. It catalyzes ubiquitination and endoplasmic reticulum-associated degradation of specific proteins.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 267 Q9UKV5

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

GP78; RNF45; AMFR

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
<u> </u>	Π	www.ahclonal.com.cn

Product Information


Purification Source Isotype Rabbit IgG Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using AMFR Rabbit pAb (A3731) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (ASO14) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.