

# NDUFS3 Rabbit pAb

Catalog No.: A3983

## Basic Information

### Observed MW

26kDa

### Calculated MW

30kDa

### Category

Primary antibody

### Applications

ELISA, WB

### Cross-Reactivity

Human, Mouse

## Background

This gene encodes one of the iron-sulfur protein (IP) components of mitochondrial NADH:ubiquinone oxidoreductase (complex I). Mutations in this gene are associated with Leigh syndrome resulting from mitochondrial complex I deficiency.

## Recommended Dilutions

WB 1:500 - 1:1000

## Immunogen Information

### Gene ID

4722

### Swiss Prot

O75489

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 37-264 of human NDUFS3 (NP\_004542.1).

### Synonyms

CI-30; MC1DN8; NDUFS3

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of extracts of THP-1 cells, using NDUFS3 antibody (A3983).  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.



Western blot analysis of extracts of various cell lines, using NDUFS3 Rabbit pAb antibody (A3983) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.