# Rad18 Rabbit mAb

Catalog No.: A4339 Recombinant



### **Basic Information**

### **Observed MW**

70kDa/90kDa

### **Calculated MW**

56kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC1058

## **Background**

The protein encoded by this gene is highly similar to S. cerevisiae DNA damage repair protein Rad18. Yeast Rad18 functions through its interaction with Rad6, which is an ubiquitin-conjugating enzyme required for post-replication repair of damaged DNA. Similar to its yeast counterpart, this protein is able to interact with the human homolog of yeast Rad6 protein through a conserved ring-finger motif. Mutation of this motif results in defective replication of UV-damaged DNA and hypersensitivity to multiple mutagens.

### **Recommended Dilutions**

**WB** 

1:500 - 1:1000

# **Immunogen Information**

**Gene ID** 56852

Swiss Prot Q9NS91

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Rad18 (Q9NS91).

# **Synonyms**

RNF73; Rad18

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Ī	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using Rad18 Rabbit mAb (A4339) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG(H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.