Aconitase 2 (ACO2) Rabbit mAb

www.abclonal.com

ABclonal

Catalog No.: A4524 Recombinant 7 Publications

Basic Information

Observed MW

85kDa

Calculated MW

85kDa

Category

Primary antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC1072

Background

The protein encoded by this gene belongs to the aconitase/IPM isomerase family. It is an enzyme that catalyzes the interconversion of citrate to isocitrate via cis-aconitate in the second step of the TCA cycle. This protein is encoded in the nucleus and functions in the mitochondrion. It was found to be one of the mitochondrial matrix proteins that are preferentially degraded by the serine protease 15(PRSS15), also known as Lon protease, after oxidative modification.

Recommended Dilutions

WB 1:1000 - 1:6000

1:200 - 1:800 **IHC-P**

IF/ICC 1:100 - 1:400

Recommended starting **ELISA**

> concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot Q99798

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ICRD; OCA8; OPA9; ACONM; HEL-S-284; Aconitase 2 (ACO2)

Contact

	400-999-6126
\bowtie	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Product Information

Source Isotype **Purification** Rabbit IgG Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of various lysates, using Aconitase 2 (ACO2) Rabbit mAb (A4524) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.




Western blot analysis of various lysates, using Aconitase 2 (ACO2) Rabbit mAb (A4524) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.



Immunofluorescence analysis of NIH-3T3 cells using Aconitase 2 (ACO2) (ACO2) Rabbit mAb (A4524) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffinembedded Human colon tissue using Aconitase 2 (ACO2) Rabbit mAb (A4524) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse kidney tissue using Aconitase 2 (ACO2) Rabbit mAb (A4524) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of paraffinembedded Rat kidney tissue using Aconitase 2 (ACO2) Rabbit mAb (A4524) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.