# **NNT Rabbit pAb**

Catalog No.: A4561 2 Publications



## **Basic Information**

### **Observed MW**

114kDa/125kDa

### **Calculated MW**

114kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes an integral protein of the inner mitochondrial membrane. The enzyme couples hydride transfer between NAD(H) and NADP(+) to proton translocation across the inner mitochondrial membrane. Under most physiological conditions, the enzyme uses energy from the mitochondrial proton gradient to produce high concentrations of NADPH. The resulting NADPH is used for biosynthesis and in free radical detoxification.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

## **Immunogen Information**

**Gene ID** 23530

**Swiss Prot** 

Q13423

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 881-1086 of human NNT (NP\_036475.3).

### **Synonyms**

GCCD4; NNT

### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using NNT Rabbit pAb (A4561) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.