

DHX9/RNA Helicase A Rabbit mAb

Catalog No.: A4563 **Recombinant** **2 Publications**

Basic Information

Observed MW

141kDa

Calculated MW

141kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P, IF/ICC, IP

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC1033

Background

This gene encodes a member of the DEAH-containing family of RNA helicases. The encoded protein is an enzyme that catalyzes the ATP-dependent unwinding of double-stranded RNA and DNA-RNA complexes. This protein localizes to both the nucleus and the cytoplasm and functions as a transcriptional regulator. This protein may also be involved in the expression and nuclear export of retroviral RNAs. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 11 and 13.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
IP	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

Immunogen Information

Gene ID

1660

Swiss Prot

Q08211

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1100-1200 of human DHX9/RNA Helicase A (Q08211).

Synonyms

LKP; RHA; DDX9; NDH2; NDHII; DHX9/RNA Helicase A

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg DHX9/DHX9/RNA Helicase A antibody (A4563). Western blot was performed from the immunoprecipitate using DHX9/DHX9/RNA Helicase A antibody (A4563) at a dilution of 1:2000.



Western blot analysis of various lysates using DHX9/RNA Helicase A Rabbit mAb (A4563) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 1s.



Confocal imaging of U-2 OS cells using DHX9/RNA Helicase A Rabbit mAb (A4563, dilution 1:100) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x.




Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded human colon carcinoma using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded human colon using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.


Validation Data




Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded human placenta using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded mouse kidney using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded mouse liver using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded rat brain using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded rat liver using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded rat spleen using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of DHX9/RNA Helicase A in paraffin-embedded rat testis using DHX9/RNA Helicase A Rabbit mAb (A4563) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.