

Arginase 1 (ARG1) Rabbit mAb

Catalog No.: A4923 **Recombinant** **23 Publications**

Basic Information

Observed MW

40kDa

Calculated MW

35kDa

Category

Primary antibody

Applications

ELISA,WB,IF/ICC

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC1164

Background

Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exist (types I and II) which differ in their tissue distribution, subcellular localization, immunologic crossreactivity and physiologic function. The type I isoform encoded by this gene, is a cytosolic enzyme and expressed predominantly in the liver as a component of the urea cycle. Inherited deficiency of this enzyme results in arginemia, an autosomal recessive disorder characterized by hyperammonemia. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB	1:2000 - 1:6000
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

383

Swiss Prot

P05089

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Arginase 1 (ARG1) (P05089).

Synonyms

ARG1; arginase-1; Arginase 1 (ARG1)

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts of Rat liver, using Arginase 1 (Arginase 1 (ARG1)) Rabbit mAb (A4923) at 1:5000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 1s.



Immunofluorescence analysis of rat liver using Arginase 1 (ARG1) Rabbit mAb (A4923) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Confocal imaging of paraffin-embedded human liver using Arginase 1 (ARG1) Rabbit mAb (A4923, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x.



Confocal imaging of paraffin-embedded mouse liver using Arginase 1 (ARG1) Rabbit mAb (A4923, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x.