

GLP-1 Rabbit mAb

Catalog No.: A4981 **Recombinant**

Basic Information

Observed MW

Calculated MW

21kDa

Category

Primary antibody

Applications

ELISA, IHC-P, IF/ICC

Cross-Reactivity

Mouse, Rat

CloneNo number

ARC1142

Background

The protein encoded by this gene is actually a preproprotein that is cleaved into four distinct mature peptides. One of these, glucagon, is a pancreatic hormone that counteracts the glucose-lowering action of insulin by stimulating glycogenolysis and gluconeogenesis. Glucagon is a ligand for a specific G-protein linked receptor whose signalling pathway controls cell proliferation. Two of the other peptides are secreted from gut endocrine cells and promote nutrient absorption through distinct mechanisms. Finally, the fourth peptide is similar to glicentin, an active enteroglucagon.

Recommended Dilutions

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID

2641

Swiss Prot

P01275

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 70-150 of human GLP-1 (P01275).

Synonyms

GLP1; GLP2; GRPP; GLP-1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data




Immunofluorescence analysis of rat pancreas using GLP-1 Rabbit mAb (A4981) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of mouse pancreas using GLP-1 Rabbit mAb (A4981) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of GLP-1 in paraffin-embedded rat pancreas using GLP-1 Rabbit mAb (A4981) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of GLP-1 in paraffin-embedded mouse pancreas using GLP-1 Rabbit mAb (A4981) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.