CBX4 Rabbit mAb

Catalog No.: A5109 Recombinant 2 Publications



Basic Information

Observed MW

53kDa/60kDa

Calculated MW

61kDa

Category

Primary antibody

Applications

ELISA,WB,IP

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC1201

Background

Enables SUMO binding activity; SUMO ligase activity; and enzyme binding activity. Involved in negative regulation of transcription by RNA polymerase II. Located in nuclear body. Part of PRC1 complex. Implicated in hepatocellular carcinoma. Biomarker of hepatocellular carcinoma.

Recommended Dilutions

WB 1:500 - 1:1000

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

Oμg extracts of whole cells

Immunogen Information

Gene ID Swiss Prot 8535 000257

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 461-560 of human CBX4 (O00257).

Synonyms

PC2; NBP16; CBX4

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\overline{a}	ı	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at -20 $^{\circ}\text{C}.$ Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Immunoprecipitation analysis of 300 μg extracts of MCF7 cells using 3 μg CBX4 antibody (A5109). Western blot was performed from the immunoprecipitate using CBX4 antibody (A5109) at a dilution of 1:1000.



Western blot analysis of various lysates using CBX4 Rabbit mAb (A5109) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.