

# [KO Validated] CDK8 Rabbit pAb

Catalog No.: A5548 **KO Validated** **1 Publications**

## Basic Information

**Observed MW**

56kDa

**Calculated MW**

53kDa

**Category**

Primary antibody

**Applications**

ELISA,WB,IP

**Cross-Reactivity**

Human, Mouse, Rat

## Background

This gene encodes a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are known to be important regulators of cell cycle progression. This kinase and its regulatory subunit, cyclin C, are components of the Mediator transcriptional regulatory complex, involved in both transcriptional activation and repression by phosphorylation of the carboxy-terminal domain of the largest subunit of RNA polymerase II. This kinase regulates transcription by targeting the cyclin-dependent kinase 7 subunits of the general transcription initiation factor IIH, thus providing a link between the Mediator complex and the basal transcription machinery. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

**WB** 1:500 - 1:2000**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

## Immunogen Information

**Gene ID**

1024

**Swiss Prot**

P49336

**Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 365-464 of human CDK8 (NP\_001251.1).

**Synonyms**

K35; IDHBA; K8

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.


## Validation Data



Immunoprecipitation analysis of 200 µg extracts of Jurkat cells, using 3 µg CDK8 antibody (A5548). Western blot was performed from the immunoprecipitate using CDK8 antibody (A5548) at a dilution of 1:1000.



Western blot analysis of lysates from wild type (WT) and CDK8 knockout (KO) HeLa cells, using [KO Validated] CDK8 Rabbit pAb (A5548) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.



Western blot analysis of lysates from Jurkat cells, using [KO Validated] CDK8 Rabbit pAb (A5548) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.