

# ICAM-1/CD54 Rabbit pAb

Catalog No.: A5597

32 Publications

## Basic Information

### Observed MW

89kDa/92kDa

### Calculated MW

58kDa

### Category

Primary antibody

### Applications

ELISA, WB, IHC-P, IF/ICC

### Cross-Reactivity

Human, Mouse, Rat

## Background

This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor.

## Recommended Dilutions

WB	1:1000 - 1:5000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

## Immunogen Information

### Gene ID

3383

### Swiss Prot

P05362

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 28-480 of human ICAM-1/CD54 (NP\_000192.2).

### Synonyms

BB2; CD54; P3.58; ICAM-1/CD54

## Contact

☎	400-999-6126
✉	cn.market@abclonal.com.cn
🌐	www.abclonal.com.cn

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates, using ICAM-1/CD54 Rabbit pAb (A5597) at 1:3000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 0.5s.




Western blot analysis of various lysates, using ICAM-1/CD54 Rabbit pAb (A5597) at 1:3000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 0.5s.




Western blot analysis of lysates from Mouse lung, using ICAM-1/CD54 Rabbit pAb (A5597) at 1:3000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 0.5s.




Immunofluorescence analysis of L929 cells using ICAM-1/CD54 Rabbit pAb (A5597) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of ICAM-1/CD54 in paraffin-embedded mouse lung using ICAM-1/CD54 Rabbit pAb (A5597) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of ICAM-1/CD54 in paraffin-embedded mouse lung using ICAM-1/CD54 Rabbit pAb (A5597) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of ICAM-1/CD54 in paraffin-embedded rat lung using ICAM-1/CD54 Rabbit pAb (A5597) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of ICAM-1/CD54 in paraffin-embedded rat lung using ICAM-1/CD54 Rabbit pAb (A5597) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of ICAM-1/CD54 in paraffin-embedded rat spleen using ICAM-1/CD54 Rabbit pAb (A5597) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of ICAM-1/CD54 in paraffin-embedded rat spleen using ICAM-1/CD54 Rabbit pAb (A5597) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.