HTRA2 Rabbit pAb

Catalog No.: A5762



Basic Information

Observed MW

37kDa

Calculated MW

49kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse

Background

This gene encodes a serine protease. The protein has been localized in the endoplasmic reticulum and interacts with an alternatively spliced form of mitogen-activated protein kinase 14. The protein has also been localized to the mitochondria with release to the cytosol following apoptotic stimulus. The protein is thought to induce apoptosis by binding the apoptosis inhibitory protein baculoviral IAP repeat-containing 4. Nuclear localization of this protein has also been observed. Alternate splicing of this gene results in multiple transcript variants encoding different isoforms.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

 $\begin{array}{c} \textbf{ELISA} & \text{Recommended starting} \\ & \text{concentration is 1 } \mu\text{g/mL}. \end{array}$

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 27429 043464

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

OMI; MGCA8; PARK13; PRSS25; HTRA2

Contact

<u>a</u>	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{\mathfrak{S}}$	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage


Store at -20 $^{\circ}\text{C}.$ Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using HTRA2 Rabbit pAb (A5762) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).



Immunofluorescence analysis of HeLa cells using HTRA2 Rabbit pAb (A5762). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.