

TGM3 Rabbit pAb

Catalog No.: A5856

Basic Information

Observed MW

77kDa

Calculated MW

77kDa

Category

Primary antibody

Applications

ELISA, WB, IF/ICC

Cross-Reactivity

Human, Mouse

Background

Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved, they all have the same amino acid sequence at their active sites and their activity is calcium-dependent. The protein encoded by this gene consists of two polypeptide chains activated from a single precursor protein by proteolysis. The encoded protein is involved in the later stages of cell envelope formation in the epidermis and hair follicle.

Recommended Dilutions

WB	1:500 - 1:1000
IF/ICC	1:10 - 1:100

Immunogen Information

Gene ID

7053

Swiss Prot

Q08188

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of human TGM3 (NP_003236.3).

Synonyms

TGE; UHS2; TGM3

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from A549 cells using TGM3 Rabbit pAb(A5856) at 1:600 dilution.
Secondary antibody:HRP Goat Anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection:ECL Basic Kit (RM00020).
Exposure time:20s.



Immunofluorescence analysis of U2OS cells using TGM3 Rabbit pAb (A5856). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.