

TDRD3 Rabbit pAb

Catalog No.: A6000

Basic Information

Observed MW

73kDa

Calculated MW

73kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human

Background

Enables chromatin binding activity; methylated histone binding activity; and transcription coactivator activity. Predicted to be involved in chromatin organization and positive regulation of transcription, DNA-templated. Located in Golgi apparatus; cytosol; and nucleoplasm. Colocalizes with exon-exon junction complex.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

81550

Swiss Prot

Q9H7E2

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

TDRD3

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of lysates from U2OS cells, using TDRD3 Rabbit pAb (A6000) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.