Coilin Rabbit pAb

Catalog No.: A6428 2 Publications



Basic Information

Observed MW

80kDa

Calculated MW

63kDa

Category

Primary antibody

Applications

ELISA,WB,IP

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene is an integral component of Cajal bodies (also called coiled bodies). Cajal bodies are nuclear suborganelles of varying number and composition that are involved in the post-transcriptional modification of small nuclear and small nucleolar RNAs. The N-terminus of the coilin protein directs its self-oligomerization while the C-terminus influences the number of nuclear bodies assembled per cell. Differential methylation and phosphorylation of coilin likely influences its localization among nuclear bodies and the composition and assembly of Cajal bodies. This gene has pseudogenes on chromosome 4 and chromosome 14.

Recommended Dilutions

WB 1:500 - 1:2000

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

Immunogen Information

 Gene ID
 Swiss Prot

 8161
 P38432

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 297-576 of human Coilin (NP_004636.1).

Synonyms

CLN80; p80-coilin; Coilin

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\overline{a}	ı	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using Coilin Rabbit pAb (A6428) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit $\lg G$ (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.