

CCL25 Rabbit pAb

Catalog No.: A6543 **2 Publications**

Basic Information

Observed MW

Refer to Figures

Calculated MW

17kDa

Category

Primary antibody

Applications

IHC-P,ELISA

Cross-Reactivity

Mouse, Rat

Background

This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for dendritic cells, thymocytes, and activated macrophages but is inactive on peripheral blood lymphocytes and neutrophils. The product of this gene binds to chemokine receptor CCR9. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

IHC-P 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

6370

Swiss Prot

O15444

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 24-150 of human CCL25 (NP_005615.2).

Synonyms

TECK; Ckb15; SCYA25; TECKvar; Ck beta-15; CCL25

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.


Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse thymus using CCL25 Rabbit pAb (A6543) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat thymus using CCL25 Rabbit pAb (A6543) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse intestin using CCL25 Rabbit pAb (A6543) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.