# TRMT1 Rabbit pAb

Catalog No.: A7116



## **Basic Information**

#### **Observed MW**

90kDa

#### **Calculated MW**

72kDa

#### Category

Primary antibody

## **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene encodes a tRNA-modifying enzyme that acts as a dimethyltransferase, modifying a single guanine residue at position 26 of the tRNA. The encoded enzyme has both mono- and dimethylase activity when exogenously expressed, and uses S-adenosyl methionine as a methyl donor. The C-terminal region of the encoded protein has both a zinc finger motif, and an arginine/proline-rich region. Mutations in this gene have been implicated in autosomal recessive intellectual disorder (ARID). Alternative splicing results in multiple transcript variants encoding different isoforms. There is a pseudogene of this gene on the X chromosome.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Swiss Prot Gene ID** 55621 Q9NXH9

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

TRM1; MRT68; TRMT1

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Ī	www.abclonal.com.cn


## **Product Information**

**Purification** Source Isotype Rabbit IgG Affinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using TRMT1 Rabbit pAb (A7116) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.