ACADS Rabbit pAb

Catalog No.: A7230



Basic Information

Observed MW

44kDa

Calculated MW

44kDa

Category

Primary antibody

Applications

ELISA,WB,IP

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a tetrameric mitochondrial flavoprotein, which is a member of the acyl-CoA dehydrogenase family. This enzyme catalyzes the initial step of the mitochondrial fatty acid beta-oxidation pathway. Mutations in this gene have been associated with short-chain acyl-CoA dehydrogenase (SCAD) deficiency. Alternative splicing results in two variants which encode different isoforms.

Recommended Dilutions

WB 1:500 - 1:2000

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

35

Swiss Prot

P16219

Immunogen Information

Immunogen

Gene ID

Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human ACADS (NP_000008.1).

Synonyms

SCAD; ACAD3; ACADS

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
$\overline{\triangle}$	ī	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Immunoprecipitation analysis of 200 μ g extracts of MCF-7 cells, using 3 μ g ACADS antibody (A7230). Western blot was performed from the immunoprecipitate using ACADS antibody (A7230) at a dilution of 1:1000.



Western blot analysis of various lysates using ACADS Rabbit pAb (A7230) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 15s.