

# $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb

Catalog No.: A7248

55 Publications

## Basic Information

### Observed MW

42kDa

### Calculated MW

42kDa

### Category

Primary antibody

### Applications

WB,IF-P,IHC-P,ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

This gene encodes one of six different actin proteins. Actins are highly conserved proteins that are involved in cell motility, structure, integrity, and intercellular signaling. The encoded protein is a smooth muscle actin that is involved in vascular contractility and blood pressure homeostasis. Mutations in this gene cause a variety of vascular diseases, such as thoracic aortic disease, coronary artery disease, stroke, and Moyamoya disease, as well as multisystemic smooth muscle dysfunction syndrome.

## Recommended Dilutions

**WB** 1:500 - 1:1000**IF-P** 1:50 - 1:200**IHC-P** 1:20 - 1:200**ELISA** Recommended starting concentration is 1  $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

59

### Swiss Prot

P62736

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

ACTSA;  $\alpha$ -Smooth Muscle Actin (ACTA2)

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Validation Data




Western blot analysis of lysates from Mouse heart, using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at 1:1000 dilution.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25 $\mu$ g per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 10s.




Immunofluorescence analysis of paraffin-embedded Human colon CA using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of paraffin-embedded mouse large intestine using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded human colon tissue using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded human esophagus tissue using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded human liver cancer tissue using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.

## Validation Data

---



Immunohistochemistry analysis of paraffin-embedded mouse testis tissue using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded rat colon tissue using  $\alpha$ -Smooth Muscle Actin (ACTA2) Rabbit pAb (A7248) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.