

Catalog No.: A7302 1 Publications



## **Basic Information**

Observed MW 60kDa

Calculated MW 56kDa

Category Primary antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse, Rat

## Background

The protein encoded by this gene contains a RING finger motif and an FHA domain. This protein has been shown to interact with several class II ubiquitin-conjugating enzymes (E2), including UBE2E1/UBCH6, UBE2E2, and UBE2E3, and may act as an ubiquitin ligase (E3) in the ubiquitination of certain nuclear proteins. This protein is also known to play a role in the DNA damage response and depletion of this protein causes cell growth inhibition and cell cycle arrest. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

WB	1:100 - 1:500
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID** 9025 **Swiss Prot** 076064

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 230-485 of human RNF8 (NP\_003949.1).

Synonyms hRNF8; RNF8

## Contact


# a 400-999-6126 x cn.market@abclonal.com.cn y www.abclonal.com.cn

# **Product Information**

**Source** Rabbit **Isotype** IgG Purification Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates, using RNF8 Rabbit pAb (A7302) at 1:400 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 90s.