

Catalog No.: A7314 8 Publications



Basic Information

Observed MW 21kDa

Calculated MW 21kDa

Category Primary antibody

Applications ELISA,WB,IF/ICC

Cross-Reactivity Human, Mouse

Background

This gene encodes a basic helix-loop-helix (bHLH) transcription factor that plays an important role in embryonic development. The encoded protein forms both homodimers and heterodimers that bind to DNA E box sequences and regulate the transcription of genes involved in cranial suture closure during skull development. This protein may also regulate neural tube closure, limb development and brown fat metabolism. This gene is hypermethylated and overexpressed in multiple human cancers, and the encoded protein promotes tumor cell invasion and metastasis, as well as metastatic recurrence. Mutations in this gene cause Saethre-Chotzen syndrome in human patients, which is characterized by craniosynostosis, ptosis and hypertelorism.

Recommended Dilutions

Immunogen Information

WB	1:100 - 1:500	Gene ID	Swiss Prot
IF/ICC	1:50 - 1:200	7291	Q15672

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 51-150 of human TWIST1 (NP_000465.1).

Synonyms

CRS; CSO; SCS; ACS3; CRS1; BPES2; BPES3; SWCOS; TWIST; bHLHa38; Twist

6

 \sim

€


400-999-6126 cn.market@abclonal.com.cn www.abclonal.com.cn

Product Information


Source Rabbit **lsotype** IgG Purification Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from 293T cells, using Twist Rabbit pAb (A7314) at 1:400 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.



Immunofluorescence analysis of HeLa cells using Twist Rabbit pAb (A7314) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Twist Rabbit pAb (A7314) at dilution of 1:100. Blue: DAPI for nuclear staining.