

SQSTM1/p62 Rabbit pAb

Catalog No.: A7758SP

124 Publications

Basic Information

Observed MW

62 kDa

Calculated MW

48 kDa/39 kDa

Category

Primary antibody

Applications

WB, IP, IF/ICC, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear factor kappa-B (NF- κ B) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF- κ B in response to upstream signals. Alternatively spliced transcript variants encoding either the same or different isoforms have been identified for this gene. Mutations in this gene result in sporadic and familial Paget disease of bone.

Recommended Dilutions

WB 1:2000 - 1:5000**IP** 0.5 μ g - 4 μ g antibody for
200 μ g - 400 μ g extracts of
whole cells**IF/ICC** 1:200 - 1:800**IHC-P** 1:200 - 1:800**ELISA** Recommended starting
concentration is 1 μ g/mL.
Please optimize the
concentration based on your
specific assay
requirements. For high-ratio
antibody dilutions
($\geq 1:10000$) a sequential
dilution method is strongly
recommended to ensure
measurement accuracy.

Immunogen Information

Gene ID

8878

Swiss Prot

Q13501

Immunogen

This information is considered to be commercially sensitive.

Synonyms

p60; p62; A170; DMRV; OSIL; PDB3; ZIP3; p62B; NADGP; FTDALS3; SQSTM1/p62

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide.

May contain 0.05% BSA as specified on the Certificate of Analysis.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Immunoprecipitation of SQSTM1/p62 from 300 µg extracts of HeLa cells was performed using 0.5 µg of SQSTM1/p62 Rabbit pAb (A7758SP). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using SQSTM1/p62 Rabbit pAb (A7758SP) at a dilution of 1:10000.




Western blot analysis of various lysates using SQSTM1/p62 Rabbit pAb (A7758SP) at 1:2000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10 s.



Confocal imaging of A549 cells using SQSTM1/p62 Rabbit pAb (A7758SP, dilution 1:400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of NIH/3T3 cells using SQSTM1/p62 Rabbit pAb (A7758SP, dilution 1:400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of PC-12 cells using SQSTM1/p62 Rabbit pAb (A7758SP, dilution 1:400) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.


Validation Data




Immunohistochemistry analysis of paraffin-embedded Human brain tissue using SQSTM1/p62 Rabbit pAb (A7758SP) at a dilution of 1:700 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using SQSTM1/p62 Rabbit pAb (A7758SP) at a dilution of 1:700 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using SQSTM1/p62 Rabbit pAb (A7758SP) at a dilution of 1:700 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human lung squamous carcinoma tissue tissue using SQSTM1/p62 Rabbit pAb (A7758SP) at a dilution of 1:700 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.