# **ACAD9 Rabbit pAb**

Catalog No.: A7798 1 Publications



### **Basic Information**

#### **Observed MW**

69kDa

#### **Calculated MW**

69kDa

#### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

### **Background**

This gene encodes a member of the acyl-CoA dehydrogenase family. Members of this family of proteins localize to the mitochondria and catalyze the rate-limiting step in the beta-oxidation of fatty acyl-CoA. The encoded protein is specifically active toward palmitoyl-CoA and long-chain unsaturated substrates. Mutations in this gene cause acyl-CoA dehydrogenase family member type 9 deficiency. Alternate splicing results in multiple transcript variants.

# **Recommended Dilutions**

**WB** 

1:500 - 1:2000

### **Immunogen Information**

**Gene ID** 28976

Swiss Prot Q9H845

# Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human ACAD9 (NP\_054768.2).

### **Synonyms**

NPD002; MC1DN20; ACAD9

### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using ACAD9 Rabbit pAb (A7798) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.