# **UCHL3 Rabbit pAb**

Catalog No.: A8156



### **Basic Information**

#### **Observed MW**

26kDa

### **Calculated MW**

26kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is a member of the deubiquitinating enzyme family. Members of this family are proteases that catalyze the removal of ubiquitin from polypeptides and are divided into five classes, depending on the mechanism of catalysis. This protein may hydrolyze the ubiquitinyl-N-epsilon amide bond of ubiquitinated proteins to regenerate ubiquitin for another catalytic cycle. Alternative splicing results in multiple transcript variants that encode different protein isoforms.

# **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

Gene ID 7347 **Swiss Prot** 

P15374

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human UCHL3 (NP\_005993.1).

## **Synonyms**

UCH-L3; UCHL3

### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using UCHL3 antibody (A8156) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.