

TMBIM1 Rabbit pAb

Catalog No.: A8249

1 Publications

Basic Information

Observed MW

35kDa

Calculated MW

35kDa

Category

Primary antibody

Applications

ELISA, WB, IF/ICC

Cross-Reactivity

Human, Mouse

Background

Enables death receptor binding activity. Involved in negative regulation of Fas signaling pathway; negative regulation of extrinsic apoptotic signaling pathway via death domain receptors; and negative regulation of protein localization to plasma membrane. Located in Golgi apparatus; endosome membrane; and lysosomal membrane.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID

64114

Swiss Prot

Q969X1

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human TMBIM1 (NP_071435.2).

Synonyms

LFG3; RECS1; MST100; PP1201; MSTP100; TMBIM1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of lysates from U-251MG cells, using TMBIM1 Rabbit pAb (A8249) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 3min.



Immunofluorescence analysis of NIH-3T3 cells using TMBIM1 Rabbit pAb (A8249) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.