MAP1LC3C Rabbit pAb

Catalog No.: A8295 1 Publications



Basic Information

Observed MW

17kDa

Calculated MW

17kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Autophagy is a highly regulated bulk degradation process that plays an important role in cellular maintenance and development. MAP1LC3C is an ortholog of the yeast autophagosome protein Atg8 (He et al., 2003 [PubMed 12740394]).

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:100

ELISA Recommended starting concentration is 1 μ g/mL. Please optimize the

concentration based on your specific assay requirements.

Immunogen Information

Gene IDSwiss Prot
440738
Q9BXW4

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

LC3C; ATG8J; MAP1LC3C

Contact

| <u>a</u> | | 400-999-6126 |
|----------------|---|---------------------------|
| \bowtie | | cn.market@abclonal.com.cn |
| \overline{a} | Т | www.ahclonal.com.cn |

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20 $^{\circ}\text{C}.$ Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using MAP1LC3C Rabbit pAb (A8295) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.