# **ZMYND8 Rabbit pAb**

Catalog No.: A8737



### **Basic Information**

#### **Observed MW**

165kDa

#### **Calculated MW**

132kDa

#### Category

Primary antibody

#### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human

### **Background**

The protein encoded by this gene is a receptor for activated C-kinase (RACK) protein. The encoded protein has been shown to bind in vitro to activated protein kinase C beta I. In addition, this protein is a cutaneous T-cell lymphoma-associated antigen. Finally, the protein contains a bromodomain and two zinc fingers, and is thought to be a transcriptional regulator. Multiple transcript variants encoding several different isoforms have been found for this gene.

### **Recommended Dilutions**

**WB** 

1:1000 - 1:2000

### **Immunogen Information**

**Gene ID** 23613

**Swiss Prot** 

Q9ULU4

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 959-1188 of human ZMYND8 (NP\_898868.1).

### **Synonyms**

RACK7; PRKCBP1; PRO2893; ZMYND8

### **Contact**

<b>a</b>	400-999-6126
<b>×</b>	cn.market@abclonal.com.cn
$\overline{\Box}$	www.ahclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of extracts of various cell lines, using ZMYND8 antibody (A8737) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.