Leader in Biomolecular Solutions for Life Science

NME1/NM23A Rabbit mAb

Catalog No.: A8802 Recombinant 1 Publications



Basic Information

Observed MW 16 kDa/18 kDa

Calculated MW 17 kDa

Category Primary antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse, Rat

CloneNo number ARC1309

Background

This gene (NME1) was identified because of its reduced mRNA transcript levels in highly metastatic cells. Nucleoside diphosphate kinase (NDK) exists as a hexamer composed of 'A' (encoded by this gene) and 'B' (encoded by NME2) isoforms. Mutations in this gene have been identified in aggressive neuroblastomas. Two transcript variants encoding different isoforms have been found for this gene. Co-transcription of this gene and the neighboring downstream gene (NME2) generates naturally-occurring transcripts (NME1-NME2), which encodes a fusion protein comprised of sequence sharing identity with each individual gene product.

Recommended Dilutions

0 - 1:2000
ed starting 5 1 μg/mL. timize the based on cific assay uirements.

Immunogen Information

Gene ID 4830

Swiss Prot P15531

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

NB; AWD; NBS; GAAD; NDKA; NM23; NDPKA; NDPK-A; NM23-H1; NME1/NM23A

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Product Information


Source Rabbit

Isotype lgG

Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using NME1/NME1/NM23A Rabbit mAb (A8802) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.