# **GNAI1** Rabbit pAb

Catalog No.: A8844



## **Basic Information**

#### **Observed MW**

40kDa

## **Calculated MW**

40kDa

#### Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Guanine nucleotide binding proteins are heterotrimeric signal-transducing molecules consisting of alpha, beta, and gamma subunits. The alpha subunit binds guanine nucleotide, can hydrolyze GTP, and can interact with other proteins. The protein encoded by this gene represents the alpha subunit of an inhibitory complex. The encoded protein is part of a complex that responds to beta-adrenergic signals by inhibiting adenylate cyclase. Two transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

Gene ID 2770 **Swiss Prot** 

P63096

## **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 50-150 of human GNAl1 ( $NP_002060.4$ ).

# **Synonyms**

Gi; HG1B; NEDHISB; GNAI1

## **Contact**

<b>a</b>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using GNAI1 Rabbit pAb (A8844) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.