

# [KO Validated] HDAC8 Rabbit mAb

Catalog No.: A8865

KO Validated

Recombinant

5 Publications

## Basic Information

### Observed MW

42kDa

### Calculated MW

42kDa

### Category

Primary antibody

### Applications

IHC-P, IF/ICC, IP, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### CloneNo number

ARC1331

## Background

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene.

## Recommended Dilutions

IHC-P 1:100 - 1:1000

IF/ICC 1:100 - 1:1000

IP 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cellsELISA Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Immunogen Information

### Gene ID

55869

### Swiss Prot

Q9BY41

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

HD8; WTS; RPD3; CDA07; CDLS5; KDAC8; MRXS6; HDACL1; C8

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


## Validation Data




Immunoprecipitation analysis of 300 µg extracts from HeLa cells using 3 µg HDAC8 antibody (A8865). Western blot was performed from the immunoprecipitate using HDAC8 antibody (A8865) at a dilution of 1:1000.



Confocal imaging of HeLa cells using [KO Validated] HDAC8 Rabbit mAb (A8865, dilution 1:100) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x.



Immunofluorescence analysis of NIH/3T3 cells using [KO Validated] HDAC8 Rabbit mAb (A8865) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain using HDAC8 Rabbit mAb (A8865) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.