# VBP1 Rabbit pAb

Catalog No.: A8990



## **Basic Information**

### **Observed MW**

23kDa

### **Calculated MW**

23kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

## **Background**

The protein encoded by this gene interacts with the Von Hippel-Lindau protein to form an intracellular complex. The encoded protein functions as a chaperone protein, and may play a role in the transport of the Von Hippel-Lindau protein from the perinuclear granules to the nucleus or cytoplasm. Alternative splicing and the use of alternate transcription start sites results in multiple transcript variants encoding different protein isoforms.

## **Recommended Dilutions**

**WB** 

1:1000 - 1:2000

## **Immunogen Information**

Gene ID 7411 **Swiss Prot** 

P61758

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-197 of human VBP1 (NP\_003363.1).

### **Synonyms**

PFD3; PFDN3; VBP-1; HIBBJ46; VBP1

### **Contact**

2		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using VBP1 Rabbit pAb (A8990) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.