Leader in Biomolecular Solutions for Life Science



Catalog No.: A9195 Recombinant 1 Publications



Basic Information

Observed MW 19kDa

Calculated MW 21kDa

Category Primary antibody

Applications ELISA,WB,IF/ICC

Cross-Reactivity Human, Mouse, Rat

CloneNo number ARC1472

Background

ADP-ribosylation factor 1 (ARF1) is a member of the human ARF gene family. The family members encode small guanine nucleotide-binding proteins that stimulate the ADPribosyltransferase activity of cholera toxin and play a role in vesicular trafficking as activators of phospholipase D. The gene products, including 6 ARF proteins and 11 ARF-like proteins, constitute a family of the RAS superfamily. The ARF proteins are categorized as class I (ARF1, ARF2 and ARF3), class II (ARF4 and ARF5) and class III (ARF6), and members of each class share a common gene organization. The ARF1 protein is localized to the Golgi apparatus and has a central role in intra-Golgi transport. Multiple alternatively spliced transcript variants encoding the same protein have been found for this gene.

Recommended Dilutions

Immunogen Information

WB	1:500 - 1:2000	Gene ID	Swiss Prot
IF/ICC	1:50 - 1:200	375	P84077

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 70-150 of human ARF1 (P84077).

Synonyms

PVNH8; ARF1

400-999-6126 6 cn.market@abclonal.com.cn \sim € www.abclonal.com.cn

Product Information

Source Rabbit


Isotype lgG

Purification Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using ARF1 Rabbit mAb (A9195) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3min.



Immunofluorescence analysis of NIH-3T3 cells using ARF1 Rabbit mAb (A9195) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.