

# THRAP3 Rabbit pAb

Catalog No.: A9396

1 Publications

## Basic Information

### Observed MW

150kDa

### Calculated MW

109kDa

### Category

Primary antibody

### Applications

ELISA, WB, IHC-P

### Cross-Reactivity

Human, Mouse, Rat

## Background

Enables phosphoprotein binding activity; thyroid hormone receptor binding activity; and transcription coactivator activity. Involved in nuclear-transcribed mRNA catabolic process; positive regulation of circadian rhythm; and regulation of RNA metabolic process. Acts upstream of or within positive regulation of transcription by RNA polymerase II. Located in nuclear speck. Part of mediator complex. Colocalizes with exon-exon junction complex.

## Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

## Immunogen Information

### Gene ID

9967

### Swiss Prot

Q9Y2W1

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 340-500 of human THRAP3 (NP\_005110.2).

### Synonyms

BCLAF2; TRAP150; THRAP3

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


## Validation Data




Western blot analysis of lysates from HeLa cells, using THRAP3 Rabbit pAb (A9396) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Immunohistochemistry analysis of THRAP3 in paraffin-embedded human brain tissue using THRAP3 Rabbit pAb (A9396) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of THRAP3 in paraffin-embedded mouse brain tissue using THRAP3 Rabbit pAb (A9396) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of THRAP3 in paraffin-embedded rat brain tissue using THRAP3 Rabbit pAb (A9396) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.