# **HECTD1** Rabbit pAb

Catalog No.: A9433



### **Basic Information**

#### **Observed MW**

300kDa

### **Calculated MW**

289kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human

## **Background**

Predicted to enable ubiquitin protein ligase activity. Predicted to be involved in anatomical structure development; positive regulation of proteasomal ubiquitin-dependent protein catabolic process; and protein K63-linked ubiquitination. Predicted to act upstream of or within several processes, including animal organ development; negative regulation of protein localization to plasma membrane; and protein autoubiquitination.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

## **Immunogen Information**

**Gene ID** 25831

Swiss Prot Q9ULT8

# Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 2518-2610 of human HECTD1 (NP\_056197.3).

## **Synonyms**

EULIR; HECTD1

### **Contact**

<b>a</b>	400-999-6126
<b>×</b>	cn.market@abclonal.com.cn
$\overline{\Box}$	www abclonal com cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using HECTD1 Rabbit pAb (A9433) at 1:3000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.